

Supporting Information

Unrevealing the Effect of Anthraquinone Metal Salts as Wide-Range Plateau Catalyst to Enhance the Combustion Properties of Solid Propellants

Ying-Lei Wang^a, Qi-Long Yan^b, Ting An^a, Bin Chen^a, Yue-Ping Ji^a, Wei Wang^a, Feng-Qi Zhao^{*a}

(^a Xi'an Modern Chemistry Research Institute, Xi'an, Shanxi 710065, China. ^b Science and Technology on Combustion, Thermal Structure, and Internal Flow Laboratory, Northwestern Polytechnical University, Xian, 710072, China.)

* To whom correspondence should be addressed: wyl-204@163.com

Contents of Supplementary

1. Characterization of AMS	3
2. The data of burning rate (u) of DB propellants with different catalysts	4
Table S1 The burning rate of DB propellants with different catalysts	4
3. The data of burning rate (u) of RDX-CMDB propellants with different catalysts	5
Table S2 The burning rate of DB propellants with different catalysts	5
4. The data of catalysis property camparation previous and our advanced catalysts for RDX- CMDB propellants	5
Table 3 The catalysis property camparation previous and our advanced catalysts for RDX- CMDB propellants	5
5. Combustion flame structures of RDX-CMDB propellants with different catalysts	6
Figure S1. Combustion flame structures of RDX-CMDB propellants with DHAAPb	6
Figure S2. Combustion flame structures of RDX-CMDB propellants with DHAAPb/DHAACu. 6	
Figure S3. Combustion flame structures of RDX-CMDB propellants with DHDNEPb.....	7
Figure S4. Combustion flame structures of RDX-CMDB propellants with DHDNEPb/DHDNECu.....	7
Figure S5. Combustion flame structures of RDX-CMDB propellants with TOCKPb.....	7

1. Characterization of AMS

(1) 1, 8-dihydroxyanthraquinone lead (**DHAAPb**): “Black powder; yield, 4.21g (94.6%); IR (KBr, ν , cm^{-1}), 3455, 1640, 1584, 1520, 1190, 981, 821, 749, 636; Anal. Found, C 37.73, H 1.349; Calcd., C 37.75, H 1.348. XRF Found, Pb 46.49, Calcd.: Pb 46.52.”

(2) 1, 8-dihydroxyanthraquinone copper (**DHAACu**): “Red powder; yield, 5.90g (97.8%); IR (KBr, ν , cm^{-1}), 3450, 1658, 1579, 1530, 1196, 978, 823, 746, 633; Anal. Found, C 49.76, H 2.965; Calcd., C 49.78, H 2.963; XRF Found, Cu 20.98, Calcd. , Cu 21.06.”

(3) 1, 8-dihydroxy-4, 5-dinitroanthraquinone lead (**DHDNEPb**): “Brown powder; yield, 5.10 g (95.3%). IR (KBr, ν , cm^{-1}): 3062, 1704, 1608, 1520, 1431, 1335, 1287, 1214, 990, 837, 693. Anal. Found, C 31.38, H 0.750, N 5.21; Calcd, C 31.40, H 0.748, N 5.23; XRF Found, Pb 38.68, Calcd.: Pb 38.70.”

(4) 1, 8-dihydroxy-4, 5-dinitroanthraquinone copper (**DHDNECu**): “Red powder; yield, 3.71 g (94.5%); IR (KBr, ν , cm^{-1}), 3055, 1678, 1598, 1504, 1428, 1331, 1289, 1210, 987, 840, 685; Anal. Found, C 42.90, H 1.024, N 7.14; Calcd. C 42.91, H 1.022, N 7.15; XRF Found, Cu 16.20, Calcd., Cu 16.22.”

(5) 1, 4, 5, 8-tetrahydroxyanthraquinone lead (**TOCKPb**): “Black powder; yield, 6.72 g (97.4%); IR (KBr, ν , cm^{-1}), 3044, 1561, 1509, 1176, 968, 802, 734, 628; Anal. Found, C 23.98, H 0.858; Calcd., C 24.00, H 0.857; XRF Found, Pb 59.66, Calcd., Pb 60.70.”

(6) 1, 4, 5, 8-tetrahydroxyanthraquinone copper (**TOCKCu**): “Red powder; yield, 3.72 g (94.2%); IR (KBr, ν , cm^{-1}), 3040, 1565, 1507, 1180, 972, 810, 738, 632; Anal. Found, C 42.51, H 1.014; Calcd., C 42.53, H 1.013; XRF Found, Cu 32.11, Calcd., Cu 32.15.”

2. The data of burning rate (u) of DB propellants with different catalysts

Table S1 The burning rate of DB propellants with different catalysts

No.	The u ($\text{mm}\cdot\text{s}^{-1}$) of DB propellants with different catalysts at different pressure(MPa)								
	2	4	6	8	10	12	14	16	18
Control 1	2.15	3.59	5.2	6.49	7.81	8.99	9.77	10.38	11.22
DB01	3.20	5.52	7.72	9.48	11.32	10.90	11.06	11.85	12.89
DB02	2.82	3.76	5.17	6.50	8.16	9.03	10.01	10.67	11.11
DB03	3.45	7.13	10.17	11.79	14.12	14.40	14.69	14.60	15.06
DB04	3.02	4.29	5.80	7.59	9.24	10.05	11.41	12.29	13.02
DB05	6.42	10.61	12.76	13.81	14.92	14.73	14.81	14.23	14.43

3. The data of burning rate (u) of RDX-CMDB propellants with different catalysts

Table S2 The burning rate of DB propellants with different catalysts

The u ($\text{mm}\cdot\text{s}^{-1}$) of RDX-CMDB propellants with different catalysts at different pressure(MPa)									
No.	2	4	6	8	10	12	14	16	18
Control 2	3.09	5.34	7.42	9.85	11.88	14.04	15.75	17.54	19.23
CMDB01	7.45	11.03	12.82	13.63	14.12	14.69	15.39	16.70	17.60
CMDB02	8.47	13.08	15.02	15.61	15.52	15.40	15.79	17.12	17.93
CMDB03	6.27	10.50	13.30	14.61	15.45	15.88	16.70	18.21	19.36
CMDB04	6.44	11.34	14.52	16.10	16.65	16.62	17.06	18.34	19.44
CMDB05	9.53	14.05	16.20	17.50	18.45	18.96	19.50	20.41	21.10
CMDB06	10.02	14.84	17.33	19.05	20.18	21.01	21.6	22.32	22.86

4. The data of catalysis property comparison previous and our advanced catalysts for RDX-CMDB propellants

Table 3 The catalysis property comparison previous and our advanced catalysts for RDX-CMDB propellants

Catalysts	Plateau region (MPa)	Width of plateau (MPa)	Burning rate on plateau ($\text{mm}\cdot\text{s}^{-1}$)
ϕ -Pb	4-10	6	7.90
ϕ -Pb/ β -Cu	6-12	6	10.2
NP/AC	4-8	4	9.00
nNP/nAC	6-12	6	13.73
PbGal	2-8	6	7.11
BTATZPb	4-12	8	9.70
DNPPb	6-10	4	10.97
TOCKPb	6-18	12	18.87
TOCKPb/TOCKCu	6-18	12	20.62

5. Combustion flame structures of RDX-CMDB propellants with different catalysts




Figure S1. Combustion flame structures of RDX-CMDB propellants with DHAAPb




Figure S2. Combustion flame structures of RDX-CMDB propellants with DHAAPb/DHAACu




Figure S3. Combustion flame structures of RDX-CMDB propellants with DHDNEPb




Figure S4. Combustion flame structures of RDX-CMDB propellants with DHDNEPb/DHDNECu




Figure S5. Combustion flame structures of RDX-CMDB propellants with TOCKPb